



**406
ARTHRITIS
CLINIC**

Get back to your 406 life!

INFUSION SUITE		PROLASTIN-C INFUSION ORDERS	
406 Arthritis Clinic LLC 2409 Arnold Ln Ste 9 Billings, MT 59102		Phone: 406-345-0211	Fax: 747-205-0742
PATIENT INFORMATION - Include Patient Demographics and Insurance Cards			
Name:		DOB:	
MEDICAL INFORMATION			
ICD10:		Patient Height:	
Patient Weight (kg):		Allergies:	
*Weigh patient prior to each infusion			
REQUIRED TESTING			
<input type="checkbox"/> Serum IgA Level: _____			
Additional labs:			
<input type="checkbox"/> Insert IV	<input type="checkbox"/> Access Port/PICC		
PREMEDICATIONS 30 minutes prior to starting			
<input type="checkbox"/> Acetaminophen:	<input type="checkbox"/> 325mg PO X1 <input type="checkbox"/> 500mg PO X1 <input type="checkbox"/> 650mg PO X1		
<input type="checkbox"/> Diphenhydramine:	<input type="checkbox"/> 25mg IVP X1 <input type="checkbox"/> 25mg PO X1 <input type="checkbox"/> 50mg IVP X1 <input type="checkbox"/> 50mg PO X1		
<input type="checkbox"/> Solumedrol:	<input type="checkbox"/> 40mg IV X1 <input type="checkbox"/> 100mg IV X1 <input type="checkbox"/> 125mg IV X1		
<input type="checkbox"/> Antihistamine:	<input type="checkbox"/> Cetirizine 10mg PO X1 <input type="checkbox"/> Loratadine 10mg PO X1		
<input type="checkbox"/> Additional PRN:			
PROLASTIN-C ORDERS			
<input type="checkbox"/> Prolastin-C	60mg/kg IV once weekly X _____		
POST INFUSION			
<input type="checkbox"/> Flush IV line with NS. D/C IV.			
<input type="checkbox"/> Flush port with 5-10mL 0.9% NS. Lock port with Heparin 10u/mL IVP (based of size port). De-access port.			
<input type="checkbox"/> Discharge home			
Referring Provider Printed:			
Referring Provider Signature:		Date:	
Referring Provider Phone:		Referring Provider Fax:	
406 Provider Printed:			
406 Provider Signature:		Date:	

**Credentials must be included*

Infusion Directions:

- Remove vial and allow to come to room temp before administration
- Pool Prolastin-C to get intended mg/kg body weight dose into an empty sterile intravenous container.
- Discard and document any drug waste
- Infuse at 0.8 ml/kg/min (approximately 15 minutes)
- Infuse using a 5-to-15-micron infusion filter

Dosing calculator: <https://www.prolastin.com/en/hcp/dosing-and-administration>