



**406
ARTHRITIS
CLINIC**

Get back to your 406 life!

INFUSION SUITE		KRYSTEXXA INFUSION ORDERS	
406 Arthritis Clinic LLC 2409 Arnold Ln Ste 9 Billings, MT 59102		Phone: 406-345-0211	Fax: 747-205-0742
PATIENT INFORMATION - If Outside Referral, Include Patient Demographics and Insurance Cards			
Name:		DOB:	
MEDICAL INFORMATION			
ICD10:		Patient Height:	
Patient Weight (kg):		Allergies:	
*Weigh patient prior to each infusion			
REQUIRED TESTING			
<input type="checkbox"/> Serum Uric Acid level: _____ (Prior to each infusion) <input type="checkbox"/> G6PD level prior to initiation: _____			
Additional labs:			
<input type="checkbox"/> Insert IV	<input type="checkbox"/> Access Port/PICC		
PREMEDICATIONS 30 minutes prior to starting			
<input type="checkbox"/> Acetaminophen:	<input type="checkbox"/> 325mg PO X1 <input type="checkbox"/> 500mg PO X1 <input type="checkbox"/> 650mg PO X1		
<input type="checkbox"/> Diphenhydramine:	<input type="checkbox"/> 25mg IVP X1 <input type="checkbox"/> 25mg PO X1 <input type="checkbox"/> 50mg IVP X1 <input type="checkbox"/> 50mg PO X1		
<input type="checkbox"/> Solumedrol:	<input type="checkbox"/> 40mg IV X1 <input type="checkbox"/> 100mg IV X1 <input type="checkbox"/> 125mg IV X1		
<input type="checkbox"/> Antihistamine:	<input type="checkbox"/> Cetirizine 10mg PO X1 <input type="checkbox"/> Loratadine 10mg PO X1		
<input type="checkbox"/> Additional PRN:			
*Diphenhydramine and Solumedrol are required per the PI			
KRYSTEXXA ORDERS			
<input type="checkbox"/> Krystexxa 8mg IV	over 2 hours every 2 weeks X _____		
POST INFUSION			
<input type="checkbox"/> Flush IV line with NS. D/C IV.			
<input type="checkbox"/> Flush port with 5-10mL 0.9% NS. Lock port with Heparin 10u/mL IVP (based of size port). De-access port.			
<input type="checkbox"/> Discharge home after 1 hour observation time			
<input type="checkbox"/> Discharge home without 1 hour observation time			
Referring Provider Printed:			
Referring Provider Signature:		Date:	
Referring Provider Phone:		Referring Provider Fax:	
406 Provider Printed:			
406 Provider Signature:		Date:	

**Credentials must be included*

Infusion Directions:

- Remove vial and allow to warm to room temperature
- Withdraw 1mL of Krystexxa from the vial and inject into a 250mL bag of NS
- Gently invert to mix. Document and discard any waste
- Infuse over no less than 2 hours, no filter tubing required

Nursing Considerations:

- Confirm patient has discontinued oral urate lowering agents