



**406  
ARTHRITIS  
CLINIC**

Get back to your 406 life!

INFUSION SUITE		TYSABRI INFUSION ORDERS	
406 Arthritis Clinic LLC 2409 Arnold Ln Ste 9 Billings, MT 59102		Phone: 406-345-0211	Fax: 747-205-0742
<b>PATIENT INFORMATION - Include Patient Demographics and Insurance Cards</b>			
Name:		DOB:	
<b>MEDICAL INFORMATION</b>			
ICD10:		Patient Height:	
Patient Weight (kg):		Allergies:	
*Weigh patient prior to each infusion			
<b>REQUIRED TESTING</b>			
<input type="checkbox"/> JCV every 3 months	<input type="checkbox"/> JCV every 6 months		
Additional labs:			
<input type="checkbox"/> Insert IV	<input type="checkbox"/> Access Port/PICC		
<b>PREMEDICATIONS 30 minutes prior to starting</b>			
<input type="checkbox"/> Acetaminophen:	<input type="checkbox"/> 325mg PO X1 <input type="checkbox"/> 500mg PO X1 <input type="checkbox"/> 650mg PO X1		
<input type="checkbox"/> Diphenhydramine:	<input type="checkbox"/> 25mg IVP X1 <input type="checkbox"/> 25mg PO X1 <input type="checkbox"/> 50mg IVP X1 <input type="checkbox"/> 50mg PO X1		
<input type="checkbox"/> Solumedrol:	<input type="checkbox"/> 40mg IV X1 <input type="checkbox"/> 100mg IV X1 <input type="checkbox"/> 125mg IV X1		
<input type="checkbox"/> Antihistamine:	<input type="checkbox"/> Cetirizine 10mg PO X1 <input type="checkbox"/> Loratadine 10mg PO X1		
<input type="checkbox"/> Additional PRN:			
<b>TYSABRI ORDERS</b>			
<input type="checkbox"/> Tysabri	300mg IV every _____ weeks X _____		
*PI requires Q4 weeks			
<b>POST INFUSION</b>			
<input type="checkbox"/> Flush IV line with NS. D/C IV.			
<input type="checkbox"/> Flush port with 5-10mL 0.9% NS. Lock port with Heparin 10u/mL IVP (based of size port). De-access port.			
<input type="checkbox"/> Discharge after 1 hour observation time			
<input type="checkbox"/> Discharge home without observation time			
Referring Provider Printed:			
Referring Provider Signature:		Date:	
Referring Provider Phone:		Referring Provider Fax:	
406 Provider Printed:			
406 Provider Signature:		Date:	

**\*Credentials must be included**

**Infusion Directions:**

- Remove vial and allow to warm to room temperature
- Withdraw 15mL of Tysabri from vial and inject into a 100mL bag of NS
- Gently invert to mix
- Infuse over 60 minutes (no filter required for tubing)

**Nursing Considerations:**

- Per the PI, patients are to be observed for the first 12 infusions. If no reactions occur, they do not have to stay for the observation hour. Defer to provider preference for observation time.